



(-)

**The Effect of Organic Fertilization in Soil Microorganisms and the
Productivity of Two Crops of Brassica Family
(Cabbage - Cauliflower)**

()

.

..

| | |
|-----------|-------------------------|
| | |
| 1 | |
| 3 | -1 |
| 4 | -2 |
| 5 | -3 |
| 9 | |
| 10 | -1-3 |
| 13 | -2-3 |
| 17 | |
| 18 | -3-3 |
| 20 | |
| 23 | |
| | -4-3 |
| | (CO ₂) -5-3 |

| | |
|--|---|
| | <p>.</p> <p>-6-3</p> <p>-7-3</p> |
| 30 | -4 |
| 31 31 32 32 32 33 33 35 35 35 35 35 36 | <p>- 5</p> <p>-1- 5</p> <p>-2-5</p> <p>-3-5</p> <p>-1-3-5</p> <p>-2-3-5</p> <p>-3-3-5</p> <p>-4-3-5</p> <p>-5-3-5</p> <p>-6-3-5</p> <p>-7-3-5</p> <p>-8-3-5</p> |

| | |
|----|------------------------|
| | -9-3-5 |
| 37 | -6 |
| 37 | |
| | -1-6 |
| | -2-6 |
| 37 | -3-6 |
| | () |
| 38 | -1-3-6 |
| 38 | -2-3-6 |
| 40 | -3-3-6 |
| 42 | -4-3-6 |
| 44 | -5-3-6 |
| 45 | -6-3-6 |
| 47 | -7-3-6 |
| 49 | -8-3-6 |
| 51 | -9-3-6 |
| 53 | -4-6 |
| | (CO ₂) () |
| | -5-6 |
| | () |
| | -6-6 |
| | () |
| 54 | |
| | -1-6-6 |

| | | |
|----|--------------------|--------|
| 56 | | -2-6-6 |
| | | -3-6-6 |
| | | -4-6-6 |
| | | -5-6-6 |
| | | -6-6-6 |
| | | -7-6-6 |
| | | -8-6-6 |
| | | -9-6-6 |
| 58 | | -7- 6 |
| 58 | (CO ₂) | |
| 59 | | |
| 61 | | -8 -6 |
| 63 | | |
| 64 | | -9-6 |
| 65 | | |
| 67 | | -10-6 |
| 69 | | |
| 70 | | |
| | | - |
| | | - |
| | | |
| | | -11-6 |
| 72 | | |

| | |
|-----|----|
| 73 | |
| 75 | |
| | - |
| 78 | - |
| 78 | |
| 79 | |
| 80 | |
| 80 | |
| 80 | |
| 82 | -7 |
| 84 | -8 |
| | -9 |
| 85 | |
| 101 | |

| | |
|------------|--|
| 103 | |
| 118 | |

:

التسميد

-) 6
- (- - -
N
)
/ N/ 50 (
)
(

Bacillus

Clostridium pasteurianum

Bacillus

.

Clostridium pasteurianum

.

.

.

.

:

.

(Liebig)
(Gilbert Laous)

C/N Ratio

.(2003)

: -2

: -3

:

% 0.1

(1988)

. ()

.(2007)

.

.

1988 (FAO)

.(1993)

794

1.663

1712 2004

(422)

.(/ 10705)

(- -)

(FAO) .

396 1988

.(1993) 205

1467 2004

(262)

.(/ 7132)

. (Lipman *et al* .,1916)

. (Dick , 1992;1994;Dilly & Blume , 1998)

(Dick , 1994;Fauci & Dick , 1994; Filip , 1998)

(García *et al.*, 1994; Valarini *et al.*, 1997;Frighetto *et al.*, 1999)

(Agrawal *et al.*, 1999)

.(2000) Mineralization

(2005)

(2001)

NH_4

CO_2

C/P C/ N

(Quelhas dos santos,2001)

CO_2

(autotrophic)

Chemoautotrophic bacteria

NH_4

Nitrosomonas Nitrobacter

.

(2007

Metcalf Eddy, 1978) NO_3

% 5-2

% 10-5

(1988

) %10-5

Pseudomonas

Clostridium Aspergillus Streptomyces

Fusarium

Bacillus Cytophaga

Micromonospora Streptomyces

Aspergillus

Penicillium

Pseudomonas

(1982)

(Postgate , 1974) *Azotobacter*

(Lynch,1983; Kennedy ,1999; 2000)

.(Insam,1990;Moreno *et al* .,1999)

.(Enwall *et al.*,2007)

(1995)

(1992)

(2003)

100 /

700-350

70-

4-3

30

(1992)

-1-3 :

(Gupta *et al* .,1988;Wong *et al* .,1999)

(Suresh *et al* ., 2004)

(Stewart *et al* .,2005)

.(Badr&Fekry,1998;Arisha&Bardisi,1999;Dauda *et al* .,2008)

microbial N

.(Zhang *et al.*, 2006)

N250 P200 K200 Kg/h

:

N200 P150 K150

N250 P200 K200

.(2003,) N200 P150 K150

()

.(Chandra *et al* ., 1993)

111

/ P / 55

/ N/

% 33

% 42

/ K 111

.% 21

.(Van Cleue and Moore,1978)

:

-2-3

(Phongran&Mosler , 2003)

(Gaur,1972;Gaur et al.,1979; Monib *et al* ,1984;Tisdale *et al* .,1985)

. (Bolan *et al* .,1987; Sanyal and De Datta,1991)

% 30-%20

% 10

% 10%20

30%

% 20 -% 10

.(Arslan *et al.*,2008) % 30

)

(2003 ,

.

.

.

()

.

.(2003)

Ammonification

Ammonifying bacteria

.(Jezierska-Tys *et al*,2008)

.

.(Mandic *et al.*, 2005)

Azotobacter

(2003)

Azotobacter

()

Azotobacter

.

(*Azotobacter*)

()

.(A)(2006)

) *Bacillus megaterium*

(

(+ + +)
 .(B)(2006)

(8)

.(Fisk *et al.*, 2001)

(2003) Edmeades

Mg Ca K P
 (Clark *et al.*.,1998)

: -3-3

(Arisha&Bardisi,1999)

.(1995)

(Zeidan *et al.*, 2001, Ahmed *et al.*,2003)

(El-Sedfy, 2003, Mahmoud *et al.* ,2004)

.(Fanous *et al.* ,2003)

(Sinaj *et al.*, 2002)

NPK :

. 70 NPK +CaCO₃ P

. ()

.(Parham *et al.*, 2003)

(NPK)

(N) 16

.(Haiyan *et al* 2006) PK

(2008) Zhong *et al*

NK NP

Long-term

()

.(Vineela *et al.*, 2008)

.(N,P,K)

.(Graham& Haynes, 2005)

NPK

.(Asit *et al.*, 2007)

(16)

()

.(Chu *et al.*, 2007)

()

Biosoild

()

/ N 200

.(Marinari *et al.*, 2000)

Marjoram

(*Azospirillum Azotobacter*)

+

.(Gewaily *et al.* ,2006)

+

+

4/1

)

(

(

)

.(Estefanous& Sawan , 2000)

NPK

(2007) Mandal *et al*

mineralizable N

.

. (Chang *et al* .,2006)

NPK

()

.(Custic *et al* .,2000)

:

-4- 3

()

.(Rao ,1982)

(1996) Hamail *et al*

(2007)El-Metwally &Abdelhamid

% 100 % 50

NPK

(7.1 4.8)

(Mahmoud *et al* .,2004)

/ (60 40)

/

)

(

.(Herencia *et al*.,2007)

: (CO₂) -5- 3

:

. (Parkin *et al* .,1996)

.(Volk,1994)

Graham& Haynes

(2005)

Biosoilde

()

(Marinari *et*

. *al.*,2000)

(2008) Gilani & Bahmanyar

Biomass

(2001) Fisk *et al*

% 30- 20

(2006) Yang Lan-Fang&Cai

% 60

.(Parkin *et al.*,1996)

% 80

7.6

(1997) Doran *et al*

.³ / 1.2

CO₂ .

20

CO₂

.(2007)

. (Prochette *et al* ., 1991)

:

) •

(CO₂

.(Parkin *et al.*,1996) () •

: -6-3

nitrite

nitrate

nitrosamines

(HNO₂)

(HNO₃)

.(Anjana, and Muhammad , 2006)

1995

(EC)

(SCF)

219)

/ 3.65 (ADI)

(60 /
. (JECFA)

/ 875
/ 500 / 900 / 1500
(Shkenan,1991)

: (2006) Fumio
.reducatase •

•
•
•

:
-
-

. reducatase
:

• -
• -

. (Addiscott *et al.*,1991)

(Ferguson *et al.* , 1990)

. (Bell *et al.* , 1990)

.(Addiscott *et al.*,1991)

(Bacher

and Lenz, 1996)

(nitrification)

(2005) Koji *et al*

% 40-20

.(Herencia *et al.*,2007)

:

-

-

(Hofman *et al.*,1989)

-

Powlson,1992)

-

.(2007 Addiscott and

-

.(Parton *et al* .,1987 , Van veen *et al.*,1981 , Lawes *et al* ., 1882)

-

.(2007)

-7-3

/ 200

/ N / 250

/ N / 300

/ N / 60

/ N / 90

/ N / 150

/ N

.(MAFF , 1994)

. (Jürgens -Gschwind , 1989)

(Rahn *et al.* 1993,

Rahn *et al.* 1998).

(Everaarts,1994)

(Batal *et al.*,1994)

(Scaife and Wurr , 1990)

(Van den Boogaard & Thorup-Kristensen , 1997)

(Blom-Zandstra,1989)

(Scaife and Turner , 1985).

(1990) Berard

. (Rahn *et al.* ,1992)

(1989) Lorenz *et al*

.(1 -)

(Lorenz *et al*

(1-)

.,1989)

| المحصول | قيمتة التسويقية العائد طن/هـ طازج | العائد الكلي طن/هـ للمحصول الطازج | امتصاص الآزوت الكلي كغ/هـ |
|--|---|--|---------------------------------|
| الكرنب | 15 | 90 | 270 |
| القرنبيط | 35 | 80 | 220 |
| الجزر | 60 | 70 | 135 |
| الكراث | 45 | 65 | 195 |
| الخس | 40 | 60 | 100 |
| البصل | 40 | 65 | 120 |
| الفجل | 30 | 25 | 60 |
| الملفوف الأبيض (قيمتة التسويقية الطازجة) | 60 | 110 | 260 |

(NH_4,NO_3)

27

. / N / 300 / N / 40 (90-0)

.

/ N / 300

. (De Neve & Hofman .,1996)

(1990) Greenwood *et al*

. (Caloin & Yu , 1984)

(1992)Weier

(- -)

. 4

.(Burns ,1996)

(Rahn *et al* .,

1992)

. 90

.(A) (Rahn *et al* .,1996)

/ 140

(Dachler,1994)

(nitrification)

Dachler

(dicyandamide)

(1994)

()

)

(

(Sady *et al* ., 1997)

.(Avrdc, 1985)

.(Sørensen, 1996)

/ N / 20

(1992) Stone and Rowse

(/ N / 240)

(B)(1996) Rahn *et al*

(Scaife & Bar Yosef , 1995)

50

. (Chamberland ,2006)

/ N / 180

(Caille *et al* ,2005; Srivastava *et al* ., 2005 ; Barcelo and Poschenrieder ,
2003 ; Clemens *et al* .,2002 ; Lombi *et al* .,2002 ; Taiz and Zeiger , 2002)
(Wintz *et al*.,2002 ; Whiting, 2001)

:

-

-

-

-

-

-

(1987) Greenwood *et al*

25

1994 (well-N) ()

(1996) Rahn *et al*

.(1996) Greenwood *et al* (N_ABLE) ()

Entec-26

.(Kołota& Adamczewska- Sowińska ,2007)

(2000) Custic *et al*

"

"

: -4

) •

(N

.

.

•

.

•

-5 :Materials and methods

-1-5 :

/

(2009-2008)(2008 – 2007)

6

:

-1

() -2

() -3

() - 4

() - 5

() -6

.((46% /N/ 50)

0 5 6×3.5 ² 21

70

50

N

(46% /N/ 50)

:

(2-)

| الكمية المضافة موسم ثاني طن / هـ | الكمية المضافة موسم أول طن / هـ | نوع السماد |
|-------------------------------------|------------------------------------|--------------|
| 13.18 | 6.35 | سماد أبقار |
| 3.18 | 6.03 | سماد أغنام |
| 3.81 | 4.6 | سماد دواجن |
| 4.44 | 4.4 | كمبوست نباتي |

(3-)

:

| نوع السماد المعدني | الكمية المضافة موسم أول طن / هـ | الكمية المضافة موسم ثاني طن / هـ |
|-----------------------|---------------------------------------|---------------------------------------|
| يوريا | 50 كغ / N هـ على شكل يوريا | 50 كغ / N هـ على شكل يوريا |
| سلفات البوتاس | K هـ على شكل سلفات البوتاس 80 كغ / | K هـ على شكل سلفات البوتاس 70 كغ / |
| سوبر فوسفات | لا يحتاج لإضافة | لا يحتاج لإضافة |

3 -2

:

(NPK)

.

-2-5 :

Brassica

Cabbage

CauliFlower

cappitata,Lizg

7

. Brassica cauliflora,Lizg

GR

.(B)%10

-3-5 :

-1-3-5 :

Olsen

660

(Olsen *et al* ., 1954)

.

.(Richards,1962) Skalar

.(Jacksan, 1958)

.(Jacksan, 1958)

: -2-3-5

10 : 1 / pH meter

EC

(Walinga *et al.*, 1995)

:

-3-3-5

(30 – 0)

)

(*Clostridium pasteurianum*

: (2006)

:Hetrotrophic bacteria () -

- 7.5- : (/) ()

30 (5) .15 - - 0.5 - 3-NaCl 1.5

(3) 10-4 ()

/) :Fungi -

30 (3) 15- - 30-Malt : (

.10⁻²

/) :

3-CaCO₃ 1-K₂HPO₄ 2- (NH₄)₂ SO₄ 15- - 10- : (

30 (5) 1-MgSO₄.7H₂O 1-NaCl

.10⁻⁴

Nystatin Agar :Actinomycetes -

-KH₂PO₄ 0.2 - 2- : (/) (Drews,1983)

(13) 15- - 0.2-MgSO₄.7H₂O 0.5
 (3) 10⁻⁴ 27-30
 : (Subba Rao , 1994) *Azotobacter* -
 -NaCl 0.2-MgSO₄ 0.2-K₂PO₄ 0.1-K₂SO₄ 20- : (/)
 (1) 20- - 5-CaCO₃ 0.2
 30 10⁻⁴

Pikovskayas Agar : -
 : (Sundara Rao and Sinah, 1963, Gaur , 1981, Pikovskaya, 1948)
 -(NH₄)₂SO₄ 5-CaPO₄ 10- 0.5 - : (/)
 (3) 10⁻⁴ 15- 0.1-MgSO₄ 0.2-KCl 0.5
 30

: -
 CaCl₂ 1- K₂HPO₄ : (/) Hechenston
 2.5 - NaNO₃ 0.01- FeCl₃ 0.1- NaCl 0.3- MgSO₄.7H₂O 0.1-
 10⁻² Dilution () 15- -

./10/ 30-28

:Spor-forming bacteria- (*Bacillus*) -

1:1
 ./5/ .10⁻⁴ ./15/ 70

:*Clostridium pasteurianum* -

.7H₂O 1-K₂HPO₄ 20- : (/) :
 20-CaCO₃ 0.5-NaCl 0.5-MgSO₄
 (MPN)

(*Clostridium Pasteurianum*)

.

: -4-3-5

.(Isermeyer, 1952,modified by Jaggi, 1976)

: -5-3-5

.(Badanie.Materialu .Roslinnego.,1972)

)

(

. (2009-2008)(2008 – 2007)

: -6-3-5

. -

: -7-3-5

:()

:

. -

-

-

-

()

-

.(2007)

: -8-3-5

. (/)

9-3-5-

:

| | | | | | | | | | | | | | |
|------------------------------|--------------|-----|--------------|-------|--------|------|--------|------|--------------|-------|--------|------|--------|
| طول التجربة بدون نطاق ٤٨,٥ م | نطاق التجربة | ٦ م | M6 | 2.5 m | M1 | 2.5m | M2 | 2.5m | M6 | 2.5 m | M1 | 2.5m | M2 |
| | | | سماد معدني | | | | | | سماد معدني | | | | |
| | | | M5 | | M6 | | M1 | | M5 | | M6 | | M1 |
| | | | كمبوست نباتي | | | | | | كمبوست نباتي | | | | |
| | | | M4 | | M5 | | M6 | | M4 | | M5 | | M6 |
| | | | سماد دواجن | | | | | | سماد دواجن | | | | |
| | | | M3 | | M4 | | M5 | | M3 | | M4 | | M5 |
| | | | سماد غنم | | | | | | سماد غنم | | | | |
| | | | M2 | | M3 | | M4 | | M2 | | M3 | | M4 |
| | | | سماد بقري | | | | | | سماد بقري | | | | |
| | | | 3.5 m | | | | | | 3.5 m | | | | |
| | | | M1 | | M2 | | M3 | | M1 | | M2 | | M3 |
| | | | شاهد | | | | | | شاهد | | | | |
| | | | مكرراً | | مكرراً | | مكرراً | | مكرراً | | مكرراً | | مكرراً |
| | | | ملف | | | | | | قرنبيط | | | | |
| عرض التجربة ٣٣,٥ م | | | | | | | | | | | | | |

:Results and discussion -6

: **-1-6**

(4-)

(4-)

| Clay | Salt | Sand | B | P | K | N | CaCO ₃ % | OM% | EC mS/cm | pH 1/10 | |
|------|------|------|------|------|-----|-------|---------------------|-----|-------------|------------|------|
| % | | | / | | | | | | | | |
| 50.9 | 14.3 | 34.8 | 0.31 | 12.1 | 207 | 63.69 | 20.3 | 1.8 | 0.45 | 8 | 0-30 |

: **-2-6**

70 (5 -)

(5-)

| K₂O % | P₂O₅ % | N % | C/N | OM % | EC mS/cm | pH 1/10 | | |
|-----------------------------------|---|----------------------|------------|-----------------------|---------------------------|--------------------------|--|--|
| 1.52 | 2.9 | 1.2 | 12/1 | 25.24 | 3.08 | 7.51 | | |
| 1.71 | 3.86 | 1.6 | 11/1 | 30.93 | 2.30 | 9 | | |
| 0.84 | 2.12 | 2.3 | 17/1 | 65.90 | 3.73 | 8.66 | | |
| 0.8 | 0.47 | 2.5 | 20/1 | 84.12 | 0.8 | 8.30 | | |
| 1.28 | 1.42 | 1.34 | 15/1 | 34.65 | 1.04 | 7.61 | | |
| 4.097 | 2.85 | 1.84 | 12/1 | 37.97 | 6.8 | 9.14 | | |
| 3.88 | 3.86 | 1.49 | 23/1 | 59.89 | 5.8 | 8.13 | | |
| 0.8 | 0.47 | 2.5 | 20/1 | 84.12 | 0.8 | 8.30 | | |

-3-6

:

: () -1-3-6

(6-)

%4.7

% 35

%32

(Parham *et al.*, 2003)

(27.3)

1 / (27.1)

1 /

.

(Enwall *et al* .,2007)

1 / (10.6)


.(Graham,Haynes,2005)

1 / (32.3)

(6 -)

(1 /)

| 24.5 ab | 22.6 a | |
|---------|--------|-----------------------------|
| 24.0 ab | 27.3 a | |
| 32.3 a | 22.7 a | |
| 18.5 bc | 20.3 a | |
| 21.6 b | 30.5 a | |
| 10.6 c | 27.1 a | |
| 10.2814 | 10.425 | LSD_(0.05) |



() (1)

: -2-3-6

(7 -)

1 / (30.3)

%53

%126

.()

1 / (29.3)

1 / (28.4)

(2003)


(4-)

(Chang *et al* .,2006)

(7 -)

(1 /)

| | | |
|----------|--------|-----------------------------|
| | | |
| 23.5 abc | 13.4 b | |
| 20.6 bc | 30.3 a | |
| 22.6 abc | 20.5 b | |
| 28.4 ab | 14.1 b | |
| 19.3 c | 17.4 b | |
| 29.3 a | 16.9 b | |
| 8.566 | 9.177 | LSD_(0.05) |



)

(2)

(

: -3-3-6

(8-)

1 / (3.3)

(2007)

1 / (9.1)

. 1 / (2.7)

1 / (10.1)

1 / (3.3)

(2003)


1 / (9.1)

1 / (4.3)

(8-)

(1 /)

| 11.4 a | 6.3 abc | |
|--------|---------|-----------------------------|
| 9.1 a | 4.3 bcd | |
| 9.8 a | 9.1 a | |
| 10.1 a | 3.3 cd | |
| 8.4 a | 6.7 ab | |
| 9.0 a | 2.7 d | |
| 5.486 | 3.038 | LSD_(0.05) |



() (3)

: -4-3-6

: (9-)

1/ (3.4)

1/ (3)

(7.2) 1 / (7.1)

Bacillus

1 /


()

.5.1 5.5 5.7 : 1 /

(9 -)

(1 /)

| | | |
|-------|--------|-----------------------|
| | | |
| 7.8 a | 1.1 c | |
| 7.1 a | 2.8 ab | |
| 7.2 a | 3.4 a | |
| 5.7 a | 1.3 c | |
| 5.5 a | 2.2 b | |
| 5.1 a | 3.0 a | |
| 2.92 | 0.65 | LSD _(0.05) |



()

(4)

: *Clostridium pasteurianum*

-5-3-6

()

(10)

Clostridium pasteurianum

0.06)

1 / (

(5 -)

(2007)


1 / (58.44)

(10 -)

1 /) *Clostridium pasteurianum*

(

| | | |
|-------|-------|--|
| | | |
| 5.13 | 9.21 | |
| 1.9 | 0.26 | |
| 57.69 | 1.97 | |
| 58.44 | 6 | |
| 0.19 | 0.06 | |
| 1.97 | 0.098 | |



) *Clostridium pasteurianum* (5)
(

: -6-3-6

(11-)

1 / (22.9)

(Mandic *et al* .,2005)

%398

%21.8 %37

(2006)

1 / (15.2)

/ (10.9) %75


1 / (10.6) 1

(Henriksson *et al* .,1976)

() (11)

(1 /)

| | | |
|---------|--------|-----------------------------|
| | | |
| 8.7 b | 4.6 e | |
| 10.9 ab | 7.4 d | |
| 9.8 b | 10.7 c | |
| 15.2 a | 17.1 b | |
| 10.6 ab | 22.9 a | |
| 6.8 b | 2.9 f | |
| 4.8 | 0.692 | LSD_(0.05) |



(6)

:

-7-3-6

(12-)

% 98 %218

(Vikram et

1 / (22.6)

pH

al.,2007)

%68

(Saha,1995)


(Debnath et al.,1994)

1 / (46.3)

(12)

(1 /)

| | | |
|--------|--------|-----------------------------|
| | | |
| 23.2 c | 7.1 cd | |
| 21.2 c | 5.5 d | |
| 22.0 c | 8.0 c | |
| 46.3 a | 22.6 a | |
| 38.0 b | 17.8 b | |
| 33.9 b | 2.3 e | |
| 7.83 | 2.47 | LSD_(0.05) |



() (7)

-8-3-6 :

(13-) 1/ (76.32)

%314

(6 -)

%169

%27

1 / (59.74)

1 / (45.02)


(2007)

(Chang *et al* .,2006)

(13)

(1 /)

| | | |
|----------|---------|-----------------------|
| | | |
| 26.50 bc | 18.42 c | |
| 20.51 bc | 49.57 b | |
| 13.68 c | 76.32 a | |
| 59.74 a | 18.67 c | |
| 45.02 ab | 16.00 c | |
| 29.82 bc | 13.16 c | |
| 26.18 | 5.98 | LSD _(0.05) |



()

(7)

: -9-3-6
(14-)

1 / (125.33)

1 / (122.67)


1 / (94.74)

(14-)

1 /)

(

| 167.54 a | 28.95 c | |
|----------|----------|-----------------------------|
| 73.50 b | 90.60 b | |
| 23.68 c | 85.09 b | |
| 102.22 b | 125.33 a | |
| 106.67 b | 122.67 a | |
| 94.74 b | 29.82 c | |
| 33.995 | 30.388 | LSD_(0.05) |



)

(9)

(

-4-6

:(CO₂)

.(15 -)

(mg CO₂g- 0.3

(mg CO₂g- 0.27

1.dm.24h-1)

1.dm.24h-1)

(mg CO₂g-1.dm.24h-1) 0.31

(0.3 mg CO₂g-1.dm.24h-1)


CO₂

(Fisk *et al* .,2007)

(15 -)

(mg CO₂g⁻¹.dm.24h⁻¹)(CO₂)

| 0.28 a | 0.25 a | |
|--------|--------|-----------------------|
| 0.31 a | 0.17 a | |
| 0.29 a | 0.2 a | |
| 0.25 a | 0.26 a | |
| 0.22 a | 0.3 a | |
| 0.3 a | 0.27 a | |
| 0.117 | 0.194 | LSD _(0.05) |



()

(10)

-5 - 6

:

(16-)

%5

3.6 10

% 54 44 33

% 6

500)

.

(/

.

()

(3-)


()

.

(16)

.(/)

| 1234.28 a | 1212.7 ab | |
|-----------|------------|-----------------------------|
| 822.93 a | 1251.86 a | |
| 685.5 a | 1089.23 b | |
| 1431.33 a | 1196.57 ab | |
| 572.76 a | 1142.03 ab | |
| 1159.58 a | 1275.42 a | |
| 1344.192 | 138.295 | LSD_(0.05) |



()

(11)

-6-6

:

: ()

-1-6-6

(17-)

1 / (28.5)

()

1 / (47.2)

()

.(Zhang *et al* .,2006)


(N,P,K)

.(Haiyan *et al* .,2006)

(17)

(1 /)

| 23.2 b | 26.5 a | |
|--------|---------|-----------------------------|
| 11.5 b | 19.7 b | |
| 13.7 b | 14.3 bc | |
| 18.4 b | 13.2 bc | |
| 22.8 b | 28.5 a | |
| 47.2 a | 12.1 c | |
| 17.76 | 6.649 | LSD_(0.05) |



()

(12)

: -2-6-6

(18 -)

1 / (29.2)

(Parham *et al.*, 2003)

/ (34.2)

1 / (34.8)

1 / (33.6)

1

1 / (31.6)

)

(2003

(Chang *et al* .,2006)

(Enwall *et al* .,2007)


(17 -)

.

(18)

(1 /)

| 15.4 c | 7.2 d | |
|---------|---------|-----------------------------|
| 23.9 bc | 12.1 cd | |
| 31.6 ab | 29.2 a | |
| 33.6 a | 7.9 d | |
| 34.2 a | 21.3 b | |
| 34.8 a | 15.8 bc | |
| 8.62 | 6.088 | LSD_(0.05) |



)

(13)

(

: -3-6-6

-)

)

(19

(

%148 1 / (10.4)

(Parham *et al.*, 2003)

.

)

1 / (11)


(2003

.(2007)

(19-)

(1 /)

| | | |
|--------|--------|-----------------------------|
| | | |
| 6.1 b | 4.2 b | |
| 7.6 ab | 6.7 b | |
| 10.8 a | 10.4 a | |
| 11.0 a | 4.5 b | |
| 9.2 ab | 4.7 b | |
| 10.7 a | 4.2 b | |
| 3.7 | 3.15 | LSD_(0.05) |



()

(14)

Bacillus

-4-6-6

:

(20-)

1 / (3.8)

1 / (4.2)


1 / (9.3)

Bacillus

(20)

(1 /)

| | | |
|--------|--------|-----------------------------|
| | | |
| 7.3 ab | 2.9 c | |
| 7.3 ab | 4.2 a | |
| 7.0 b | 2.1 d | |
| 9.3 a | 3.8 ab | |
| 9.0 ab | 2.9 c | |
| 7.1 ab | 3.4 bc | |
| 2.3 | 0.652 | LSD_(0.05) |



()

(15)

: *Clostridium pasteurianum*

-5-6-6

()

(- -) *Clostridium pasteurianum*
(21)

Clostridium pasteurianum

(2007)

(10-)

. *Clostridium pasteurianum*

(21)

1 /) *Clostridium pasteurianum*

(

| | | |
|-------|-------|--|
| | | |
| 7.2 | 0.058 | |
| 0.089 | 0.192 | |
| 2.53 | 0.192 | |
| 5.56 | 0.058 | |
| 0.183 | 0.192 | |
| 1.851 | 0.057 | |

: -6-6-6

(22-)

-)

(-

(-)


(25)

(Enwall *et al* ,2007)

(22)

(1 /)

| | | |
|---------|--------|-----------------------------|
| | | |
| 6.8 c | 6.8 ab | |
| 11.8 bc | 8.9 a | |
| 18.6 a | 8.95 a | |
| 16.0 ab | 5.8 bc | |
| 11.0 bc | 1.4 d | |
| 13.6 ab | 4.2 c | |
| 5.54 | 2.311 | LSD_(0.05) |



() (17)

: -7-6-6

(23)

1 / (36)

(Saha,1995)

.(Debnath,1994)


(-)

1 / (49.7)

(23)

(1 /)

| | | |
|---------|---------|-----------------------------|
| | | |
| 7.6 c | 12.2 b | |
| 24.2 b | 12.4 b | |
| 23.9 b | 11.3 bc | |
| 32.7 b | 12.7 bc | |
| 18.6 bc | 36 a | |
| 49.7 a | 7.2 c | |
| 14.52 | 5.102 | LSD_(0.05) |



)

(18)

(

: -8-6-6

(24-)

1 / (28.21)

()


1 / (98.73)

(2007)

(24)

(1 /)

| | | |
|---------|----------|-----------------------------|
| | | |
| 30.70 b | 3.42 c | |
| 30.95 b | 4.27 c | |
| 98.73 a | 28.21 a | |
| 37.04 b | 6.93 bc | |
| 90.24 a | 10.26 bc | |
| 16.46 c | 14.53 b | |
| 9.24 | 9.486 | LSD_(0.05) |



() (19)

:

-9-6-6

-) ()

(25

1 / (24.07)

%N

(2)

1 / (23.76)

. %P₂O₅

1 / (20.04)

(- -)

(6.24)

1 /


) 15.5 %

(

(25-)

(1 /)

| 2.06 d | 1.88 b | |
|--------|---------|-----------------------|
| 2.78 c | 20.04 a | |
| 6.24 a | 23.76 a | |
| 1.48 d | 24.07 a | |
| 5.04 b | 1.20 b | |
| 1.69 d | 1.84 b | |
| 0.59 | 7.682 | LSD _(0.05) |



(20)

()

-7-6

: (CO₂)
(26 -)

mg CO₂ g-1dm.24 h-1 (0.23)

mg CO₂ g-1dm.24 h-1 (0.2)

CO₂

(0.69)

mg CO₂ g-1dm.24 h-1 (0.74)

mg CO₂ g-1dm.24 h-1 (0.64)


mg CO₂ g-1dm.24 h-1

.(21-20-17-16) CO₂

(26)

mg CO₂ g-1dm.24 h-1(CO₂)

| | | |
|---------|---------|-----------------------------|
| | | |
| 0.596 b | 0.19 a | |
| 0.62 b | 0.2 a | |
| 0.62 b | 0.199 a | |
| 0.64 ab | 0.17 a | |
| 0.69 ab | 0.16 a | |
| 0.74 a | 0.23 a | |
| 0.12 | 0.13 | LSD_(0.05) |



() (21)

-8-6

:


(27-)

(Herencia *et al.*,2007)

(27)

.(/)

| | | |
|-------------|------------|-----------------------------|
| | | |
| 60.55 c | 961.13 a | |
| 223.67 a | 1225.182 a | |
| 127.96 abc | 1145.938 a | |
| 173.06 ab | 1098.821 a | |
| 148.995 abc | 1157.579 a | |
| 113.79 bc | 1232.113 a | |
| 102.96 | 271.12 | LSD_(0.05) |



()

(22)

(5-)

()

)

(

-9-6

:

(28-)


()

()

(28)

(/)

| 47.05 c | 43.84 a | |
|----------|---------|-----------------------------|
| 65.60 a | 42.97 a | |
| 52.11 bc | 36.60 a | |
| 62.02 ab | 44.38 a | |
| 46.16 c | 39.96 a | |
| 43.10 c | 37.61 a | |
| 12.26 | 16.9 | LSD_(0.05) |



() (23)


(29-)

()

(29-)

(/)

| | | |
|----------|-----------|-----------------------------|
| | | |
| 24.29 ab | 20.79 c | |
| 27.77 a | 30.19 a | |
| 26.37 a | 27.63 ab | |
| 24.97 ab | 30.77 a | |
| 26.57 a | abc 26.77 | |
| 19.08 b | 23.60 bc | |
| 6.004 | 6.09 | LSD_(0.05) |



() (24)

:

:

-

(30-)

()

(Chang and Salomon , 1978)

(-)

(

(Wilkinson,1979)

: (30)

%

| 1.43 c | 1.49a | |
|----------|-------|-----------------------------|
| 1.89 a | 1.58a | |
| 1.45 bc | 1.80a | |
| 1.81 ab | 1.49a | |
| 1.8 abc | 1.71a | |
| 1.61 abc | 1.76a | |
| 0.378 | 0.464 | LSD_(0.05) |

: -

(31-)

(Chang and ()

Salomon , 1978)

.

.(Sims,1995)

()

.

: (31)

%

| 0.07a | 0.08a | |
|----------|-------|-----------------------------|
| 0.09 7 a | 0.06a | |
| 0.08a | 0.07a | |
| 0.076 a | 0.06a | |
| 0.097 a | 0.07a | |
| 0.08 a | 0.08a | |
| 1.69 | 1.57 | LSD_(0.05) |

:

: -

(32-)

(-)

(Chang and Salomon , 1978, Wilkinson,1979)

Azotobacter

(Musa,1967,1974)

(Zane & Basil, 1980; Theodora *et al.*, 2003; Al-

. Tarawneh,2005)

: (32)

%

| | | |
|---------|-------|-----------------------------|
| | | |
| 1.55 ab | 1.72a | |
| 2.01 a | 1.55a | |
| 2.05 a | 1.77a | |
| 1.74 ab | 1.72a | |
| 1.9 ab | 1.72a | |
| 1.35 b | 1.64a | |
| 0.59 | 2.09 | LSD_(0.05) |

: -

.(33-)

: (33)

%

| | | |
|-------|-------|-----------------------------|
| | | |
| 0.08a | 0.08a | |
| 0.11a | 0.09a | |
| 0.1a | 0.09a | |
| 0.08a | 0.09a | |
| 0.08a | 0.09a | |
| 0.07a | 0.09a | |
| 2.49 | 1.65 | LSD_(0.05) |

| | | |
|-------|--------|----|
| | : | -7 |
|) | | |
| (46% | /N/ 50 | |
| : | | |
| | | .1 |
| . | | |
| | | .2 |
| | | |
| | . | |
| | | .3 |
| | . | |
| | | .4 |
| | | |
| (| | .5 |
|) | | |
| | . | |
| | | .6 |
| | | |
| | . | |

.7

()

.

.8

. *Bacillus*

.

.9

(-)

.10

.

.11

.

.12

.

.13

.(-) (-)

: -8

() .1

(- - -)

.

.2

.()

: -9

-1-9

:

: (34)

:

| B ppm | K ppm | P ppm | | |
|----------|--------------|----------|---------|--|
| 0.38a | 268.87a | 11.96a | | |
| 0.32a | 283.01a | 11.05a | | |
| 0.45a | 250.83a | 13.05a | | |
| 0.34a | 223.73a | 13.06a | | |
| 0.76a | 201.39a | 11.30a | | |
| 0.36a | 277.17a | 12.41a | | |
| 0.92 | 102.527 | 3.659 | LSD0.05 | |
| 0.13 ab | 105.38 c | 3.29 c | | |
| 0.15 ab | 247.79 a | 45.66 a | | |
| 0.07 b | 154.81 b | 11.51 bc | | |
| 0.21 ab | 121.17 c | 19.4 b | | |
| 0.27 a | 118.56 c | 5.2 bc | | |
| 0.08 b | 126.48 bc | 3.39 c | | |
| 0.17 | 29.78 | 14.4 | LSD0.05 | |

:

-

(34-)

(-)

(Gliessman *et al.*,1996)

(Edmeades ,2003) (Clark *et al .*,1998)

(Sanyal and De Datta,1991)

(Ruiz , 2002) (Laboski and Lamb , 2003)

H_2CO_3 CO_2

(2007 Tisdale *et al .*,1985)

(-)

-)

.(

.(Monib *et al* ,1984)

(1996,) (Maclaren and Peterson , 1967)

:-
-)

(34

(–)
 (Gliessman *et al* ..,1996;Bulluck *et al* .,2002;Edmeades , 2003)
 (Andrews *et al* .,2002)

()

.

(Saad ,1999 ; Marinari *et al* .,2000)

: -

(34-)

. (–)

:

: : (35)

:

| B ppm | K ppm | P ppm | | |
|----------|----------|----------|---------|--|
| 0.43a | 268.87a | 11.96a | | |
| 0.44a | 283.01a | 11.05a | | |
| 0.50a | 250.83a | 13.05a | | |
| 0.41a | 223.73a | 13.06a | | |
| 0.32a | 201.39a | 11.30a | | |
| 0.63a | 277.17a | 12.41a | | |
| 0.4 | 86.903 | 2.051 | LSD0.05 | |
| 0.31a | 176.68a | 12.32 b | | |
| 0.51a | 276.48a | 55.48 a | | |
| 0.45a | 147.52a | 27.3 b | | |
| 0.52a | 145.62a | 27.57 b | | |
| 0.46a | 109.04a | 11.01 b | | |
| 0.55a | 105.54a | 9.8 b | | |
| 0.28 | 385.95 | 23.79 | LSD0.05 | |

:

-

(35-)

(Bolan *et al* .,1987,Singh et Kapoor , 1999)

Gaur et al.,1979, Gaindh and Gaur

,1991,and Gaur,1972)

:() -

-)

.(

) -

:(

.(-)

) *Clostridium pasteurianum* -

:(

Clostridium pasteurianum

.

) -

:(

.

:() -

.

) -

:(

.

) -

:(

.

:() -

:() -

(36)

:2007-2008

| | | | |
|--------|----------|--------|-------------------------------------|
| | | | |
| 0.384 | -0.322 | 0.18 | |
| 0.367 | 0.13 | 0.063 | |
| 0.014 | -0.492 | -0.05 | |
| 0.181 | -0.509 | -0.515 | |
| -0.74 | 0.178 | 0.328 | <i>Clostridium pasteurianum</i> |
| -0.068 | -0.822* | -0.03 | |
| 0.862* | 2.48E-16 | 0.006 | |
| 0.458 | 0.123 | -0.766 | |
| -0.349 | -0.531 | 0.195 | |
| 0.034 | 0.42 | 0.069 | |
| -0.364 | 0.128 | -0.316 | |

:2008-2009

2008-

(37)

:2009

:() -

:() -

)

(-

:() -

-)

.(-

) -

:(

. .(- -)

-) *Clostridium pasteurianum* -

:(-

Clostridium pasteurianum

) -

:(

. (- -)

:(- -) -

| | | | | |
|----|---|----|----|---|
| | |) | | - |
| | | | :(| |
| | . | | | |
| - | |) | | - |
| | | | :(| - |
| | . | | | |
| :(| |) | | - |
| | . | | | |
| :(| - | - |) | - |
| | | . | | |
| | | .(| - | - |
| | | |) | |

(37-)

2008-2009

| | | | |
|--------|---------|--------|-------------------------------------|
| | | | |
| 0.511 | 0.052 | -0.146 | |
| -0.610 | -0.889* | -0.187 | |
| 0.564 | 0.0515 | 0.427 | |
| -0.158 | -0.631 | -0.269 | |
| -0.168 | -0.379 | 0.026 | <i>Clostridium pasteurianum</i> |
| -0.085 | -0.335 | -0.349 | |
| -0.455 | -0.362 | 0.256 | |
| -0.322 | 0.009 | 0.199 | |
| 0.45 | 0.081 | -0.253 | |
| -0.268 | -0.323 | 0.397 | |
| -0.093 | -0.777 | -0.103 | |

:2007-2008

2007-

(38-)

:2008

:() -

-)

. (-

:() -

)

.(- -

:(- -) -

-)

. (

-) -

:(-

. .(- -)

-) *Clostridium pasteurianum* -

:(-

(-) *Clostridium pasteurianum*

.

-) -

:(-

(-)

.

| | | |
|----|----|---|
| :(|) | - |
|) | | |
| . | (| - |
| - | - |) |
| | | - |
| | :(| |
| . | | |
| - |) | - |
| | :(| - |
| . | | |
| :(|) | - |
| . | | |
| :(| - | - |
| |) | - |
| . | | |

(38-)

| | | | |
|--------|--------|--------|-------------------------------------|
| | | | |
| 0.487 | 0.416 | 0.56 | |
| 0.209 | 0.317 | 0.267 | |
| -0.174 | -0.134 | 0.207 | |
| 0.375 | 0.484 | 0.162 | |
| -0.091 | 0.445 | -0.432 | <i>Clostridium pasteurianum</i> |
| 0.013 | 0.434 | 0.427 | |
| -0.81 | 0.208 | -0.37 | |
| 0.075 | -0.516 | -0.711 | |
| 0.23 | 0.147 | -0.494 | |
| 0.387 | 0.03 | 0.14 | |
| 0.603 | 0.863* | -0.19 | |

:2008-2009

2007-

(39-)

:2008

:(- -) -

:() -

)

.(- -

:(- -) -

-)

(

-) -

:(-

. .(- -)

-) *Clostridium pasteurianum* -

:(-

Clostridium pasteurianum

-) -

:(-

. (-)

:(- -) -

)

(-

| | | | |
|----|---|---|----|
| - | - |) | - |
| | | | :(|
| | | . | |
| | - |) | - |
| | | | :(|
| | | . | |
| :(| - | - |) |
| | | | - |
| | | | . |
| :(| - | - |) |
| | | | - |
| | | | . |

(39-)

.

| | | | |
|---------|--------|--------|-------------------------------------|
| | | | |
| 0.692 | 0.272 | -0.101 | |
| 0.023 | -0.213 | -0.135 | |
| 0.360 | 0.230 | 0.227 | |
| 0.056 | -0.267 | -0.188 | |
| -0.386 | -0.218 | -0.326 | <i>Clostridium pasteurianum</i> |
| 0.712 | 0.423 | 0.009 | |
| 0.121 | -0.326 | -0.143 | |
| -0.875* | -0.79 | -0.671 | |
| 0.414 | -0.316 | -0.053 | |
| -0.601 | -0.602 | -0.478 | |
| 0.595 | 0.591 | 0.807 | |

1. **2003** :
 -
2. **2001** :
 -
3. **1995** :
 -
4. **2005** :
 -
5. **1982** :
 -
6. **2003** :
 -
7. **2006** :
 -
8. **2006** :
 -
- 355 - 335 - 2 - (22) -
- (B)

| | | |
|----------|------------|----------|
| | - (2003) | .9 |
| - | | |
| | . (47) | |
| - | :(2006) | .10 |
| | | |
| | . (2007) | . 11 |
| | . (2003). | . 12 |
| | | |
| -1994 | (1) : | .13 |
| | | .1995 |
|) | - (2007) | . 15 |
| | .108-103 | .(|
| | . (2003) | .16 |
| | | |
| | : (1992) | .17 |
| () | (1993) | .18 |
| | | .19 |
| | - | : (1988) |
| : (2000) | | .20 |
| | | |

:

1. **Addiscott, T.M. Whitmore , A . P. and Powlson , D.S.(1991).** Farming , fertilizers and the nitrate problem , CABI , Walling ford .
2. **Addiscott, T.M. and Powlson , D.S. 1992.** Partitioning losses of nitrogen fertilizer between leaching and denitrification. *Journal of Agricultural Science* 118: 101-107.
3. **Agrawal, A.A.; Tuzun, S. & Bent, E. (1999).** Induced plant defenses against pathogens and herbivores: biochemistry, ecology and agriculture. St. Paul, APS Press. 390p.
4. **Ahmed, M. K. A.; Afifi, M. H.; Mohamed, M. F. (2003).** Effect of biofertilizers, chemical and organic fertilizers on growth, yield and quality of some leguminous crops. *Egypt. J. Agron.*, v. 25, n. 1, p. 45-52.
5. **Al-Tarawneh, A.A.,(2005).** The effects of two types of organic manure and NPK on growth, yield and quality of lettuce and strawberry. M. Sc.Thesis, Mu'tah University, Jordan.
6. **Anjana, Shahid Umar and Muhammad Iqbal.(2006).** Nitrate accumulation in plants, factors affecting the process, and human health implications. *Agron. Sustain. Dev.* 27 (2007) 45-57.
7. **Andrews, S.S., J.P. Mitchell, R. Mancinelli, D.L. Karlen, T.K. Hartz, W.R. Horwarth, G.S. Pettygrove, K.M. Scow, and D.S. Munk. (2002).** On-farm assessment of soil quality in California's Central Valley. *Agron. J.* 94:12–23.
8. **Arisha H.M. and A. Bradisi, (1999).** Effect of mineral fertilizers and organic fertilizers on growth, yield and quality of potato under sandy soil conditions. *Zagazig J. Agric. Res.*, 26: 391–405.
9. **Arslan. E.İşıl, Öbek. Erdal, Kirbag. Sevda, İpek. Ubeyde and TOPAL Murat (2008).** Determination of the Effect of Compost on Soil Microorganisms . *International Journal of Science & Technology*, No 2, 151- 159 ,Volume 3.
10. **Asit, Mandal; Ashok, K.Patra; Dhyan, Singh; Anand, Swarup and R.Ebhin, Masto (2007) –** Effect of long-term application of manure and fertilizer on biological and biochemical activities in soil during crop development stages, *Bioresource Technology*, vol 98(18), 3585-3592.
11. **Avrdc(1985).** Effect of time, form, and concentration of nitrogen application on Chinese cabbage tipburn. Progress Report, Asian Vegetable Research and Development Center, 385-393 Tainan, Taiwan.

12. **Badanie.Materiału Roslinnego.CZ.II.(1972)** . Metody badan laboratoryjnych w stacjach chemiczno – rolniczych . Instytut uprawy nawożenia i gleboznawstwa.
13. **Badr, L.A.A. and W.A. Fekry,(1998).** Effect of intercropping and doses of fertilization on growth and productivity of taro and cucumber plants.1-vegetative growth and chemical constituents of foliage. Zagazig J.Agric. Res., 25: 1087–101.
14. **Bacher W., and Lenz F., (1996).** Ammonium deposit fertilization to reduce nitrate leaching. Proc. Workshop Ecological Aspects Veg. Fert. (eds Hahndel, R. and Wichman, W. Acta Hort., 428: 123-130.
15. **Barcelo , J . and Poschenrieder , C . (2003)** . Phytoremediation principles and perspectives . Contribution to science , 2(3):333-344.
16. **Batal K.M., Bondari K., Granberry D.M., and Mullinix B.G. (1994).** Effects of source, rate, and frequency of N application on yield, marketable grades and rot incidence of sweet onion (*Allium cepa* L. cv. Granex-33). Journal of Horticultural Science 69: 1043-1051.
17. **Bell .S. G. and Codd. G. A.(1990).** this Issue, p. 109.
18. **Berard L.S., (1990).** Effects of nitrogen fertilization on stored cabbage. I: Development of physiological disorders on tolerant and susceptible cultivars. J. Hort Sci., 65: 289-296
19. **Bin, J.,(1983).** Utilization of green manure for raising soil fertility in China.Soil Sci., 135: 65–9.
20. **Blom-Zandstra, M.(1989).** Nitrate accumulation in vegetables and its relationship to quality. Ann. Appl. Biol., 115: 553-561
21. **Bolan N. S. , A.D. Robson, N. J. Barrow(1987).** Effects of vesicular-arbuscular mycorrhizal on the availability of iron phosphates to plants. Plant and Soil 99.401-410.
22. **Bulluck, L.R., M. Brosius, G.K. Evanylo, and J.B. Ristaino. (2002).** Organic and synthetic amendments influence soil microbial, physical and chemical properties on organic and conventional farms. Appl. Soil Ecol. 19:147–160.
23. **Burns I.G., (1996).** A Mechanistic Theory for the Relationship between Growth Rate and the Concentration of Nitrate-N or Organic-N in Young Plants derived from Nutrient Interruption Experiments. Horticulture Research International, Wellesbourne Annals of Botany 74: 159-172.

24. **Caille , N ; Zhao , F,G . and Mcgrath , S.P.(2005).** Comparison of root absorption , translocation and tolerance of arsenic in the hyperaccumulator *pteris vittata* and the non hyperaccumulator *pteris tremula* .*New phytologist* 165:755-761.
25. **Caloin M., and Yu. O. (1984).** Analysis of the time course of changes in nitrogen content of *Dactylis glomerata* L. Using a model of plant growth. *Annals of Botany* 54: 69-76.
26. **Chandra . SU . Sarath , Lee . A , Perrott . k.w , Rajan .S.S.S, Oliver and Gravett .I . M .(1993) .** Effects of phosphate fertilizer applications on microorganisms in pastoral soil . *Australian Journal of soil research.* 31(3) 299-309.
27. **Chang P-T, Salomon M. (1978).** Metals in grains sold under various labels—Organic, natural and conventional. *J FoodQual*;1:373–77.
28. **Chang .Ed-Haun, Chung .Ren-Shih and TSAI. Yuong-How.(2006).** Effect of different application rates of organic fertilizer on soil enzyme activity and microbial population. *Soil Science & Plant Nutrition .Vol 53 Issue 2, P: 132 – 140*
29. **Chamberland, E(2006).**Critical nitrate concentration in the leaves in relation to the yield and residues in a cabbage crop. *Canadian Journal of Plant Science.*
30. **Haiyan, Chu , ; Lin, Xiangui ; Fujii, Takeshi ; Morinmoto , Sho ; Yagi, Kazuyuki ; Hu, Junli and Zhang, Jiabao (2007) –** Soil microbial biomass, dehydrogenase activity, bacterial community structure in response to long-term fertilizer management, *Soil Biology and Biochemistr*,vol 39(11), 2971-2976.
31. **Clark, M.S., W.R. Horwath, C. Shennan, and K.M. Scow.**
(1998). Changes in soil chemical properties resulting from organic and low input farming practices. *Agron. J.* 90:662–671.
32. **Clemens,S;Palmgren , M. G . and Kramer , U . (2002).** Along way ahead : Understanding and engineering plant metal accumulation . *TRENDS in plant science* 7(7),309-314.
33. **Custic.M, Poljak . M , Coga.L, Cosic.T, Toth. N , Pecina.M.(2000).**The influence of organic and mineral fertilization on nutrient status , nitrate accumulation , and yield of head chicory . *Faculty of Agriculture , University of Agriculture, Zagreb , Croatia .*

34. **Dachler, M. (1994).** Stickstoffangebot und Nitratakkumulation bei Feldgemüse. (Nitrogen supply and nitrate accumulation of field vegetables). Jahresbericht 1994 des Bundesamtes und Forschungszentrums für Landwirtschaft, Wien (Annual report of the federal office and research centre of agriculture 1994, Vienna), 77-79
35. **Dauda, S.N., F.A. Ajayi and E. Ndor, (2008).** Growth and yield of water melon (*Citrullus lanatus*) as affected by poultry manure application. *J. Agric. Soc. Sci.*, 4: 121–4.
36. **Debnath , A , C . Das, and D . Mukherjee .(1994) .** Studies on the decomposition of non – conventional organic wastes in soil . *Microbiological Research* . 149 : 2 , 195 – 201
37. **De Neve S., and Hofman G., (1996).** Modelling N mineralization of vegetable crop residues during laboratory incubations. *Soil Biol. Biochem.* 28: 1451 - 1457.
38. **Dick, R.P. (1992).**A review: long-term effects of agricultural systems on soil biochemical and microbial parameters. *Agric. Ecosyst. Environ.*, 40:25-60.
39. **Dick, R.P. (1994).** Soil enzyme activities as indicators of soil quality. In: Doran, J.W.; Coleman, D.C.; Bezdicek, D.F. & Stewart, B.A., eds. *Defining soil quality for a sustainable environment*. Minneapolis, Soil Science Society of America,. p.107-124. (SSSA Special Publication, 35)
40. **Dilly, O. & Blume, H.P. ,(1998).** Indicators to assess sustainable land use with reference to soil microbiology. *Adv. GeoEcol.*,31:29-36.
41. **Doran, J.W., T. Kettler, M. Liebig, and M. Tsivou. (1997).** Solvita soil test evaluation, personal communication.
42. **Drews , G . (1983) .** Mikrobiologisches praktikum, Springer – veolago . Berlin, Heidelberg . New York . Tokyo.
43. **Edmeades, D. C.(2003).** The long-term effects of manures and fertilizers on soil productivity and quality: a review. *Nutr. Cycl. Agroecosys.*, v. 66, p. 165-180.
44. **El-Metwally, I.M.A.and Abdelhamid, M.T.(2007).** Weed control under integrated nutrient management systems in faba bean (*Vicia faba*) production in Egypt. *Planta daninha* vol.26 no.3 Viçosa 2008.
45. **EL-Sedfy, O. M. F. (2003).** Residual effect of different organic materials on some physical properties of sandy soil production and

nutrient content of Egyptian clover. J. Agric. Sci. Mansoura Univ., v. 28, n. 8, p. 6505-6515.

46. **Enwall, K .; Bertilsson , S.; Cederlund , H .; Stenstrom, J .; and Hallin, S .(2007)** : Long – term impact of fertilization on activity and composition of bacterial communities and metabolic guilds in agricultural soil Biochemistry : 106-115.

47. **Estefanous . A . N .,Sawan . O . M . (2000)** . Effect of inoculation with phosphate-bacteria , sawdust compost and nitrogen sources on Okra yield and some properties of calcareous soil . International Society for Horticultural Science .

48. **Everaarts A.P., (1994).** Nitrogen fertilization and heart rot in broccoli. Netherlands Journal of Agricultural Science, 42: 195-201.

49. **Fauci, M.F. & Dick, R.P. (1994).** Microbial biomass as an indicator of soil quality: effects of long-term management and recent soil amendments. In: Doran, J.W.; Coleman, D.C.;Bezdicek, D.F. & Stewart, B.A., eds. Defining soil quality for a sustainable environment. Minneapolis, Soil Science Society of America. 17:229-234. (SSSA Special Publication, 35).

50. **Fanous, N. E.; Abd EL-Rasoul, SH. M.; Hassan, M. M. (2003).** Sustainable peanut production through integration between bioorganic and chemical fertilizers. J. Agric. Sci. Mansoura Univ., v. 28, n. 5, p. 4243-4258.

51. **Ferguson .A. J. D., Pearson. M. J. and Reynolds. C. S.(1990).**, this Issue, p. 27.

52. **Filip, Z.K. (1998).**Soil quality assessment: an ecological attempt using microbiological and biochemical procedures. Adv. GeoEcol.,31:21-27.

53. **Fisk. C., Melany;Fahey .J, Timothy (2001)** - Microbial Biomass and Nitrogen Cycling Responses to Fertilization and Litter Removal in Young Northern Hardwood Forests, Biogeochemistry, vol52(2),201-223.

54. Food and Agriculture Organization of the United Nations ,Rome,vol 42.1988.FAO Production year book,350p.

55. **Frighetto, R.T.S.; Valarini, P.J.; Tokeshi, H. &Oliveira, D.A. (1999).** Action of effective microorganisms EM on microbial, biochemical and compaction parameters of sustainable soil in Brazil. In: International Conference On Kyusei Nature Farming, 5.,Bangkok, 1997.

Proceedings. Bangkok, Asia Pacific Natural Agriculture Network, p.159-164.

56. **Fumio,Sato.(2006).** How do nitrate ions accumulate in the head leaves of cabbage? Agriculture and Horticulture. Pub. Country;Japan. ISSN:0369-5247. VOL.81;NO.9;PAGE.989-993.

57. **Gaind S., A.C.Gaur (1991) .** Thermotolerant phosphate solubilizing microorganisms and their interaction with mung bean.Plant and Soil.137:141-149.

58. **Garcia, C.; Hernandez, T.; Costa, F. & Ceccanti, B.(1994).**Biochemical parameters in soils regenerated by the addition of organic wastes. Waste Managem. Res., 12:457-466.

59. **Gaur A. C.(1972).** Role of phosphate solubilizing microorganisms and organic matter in soil productivity. Soil Productivity: Nat. Acad. Sci. India, 259-268.

60. **Gaur A.C. , D. Arora , N. Prakash (1979).** Electron microscopy of some rock phosphate dissolving bacteria and fungi. Folio Micriobiol.24 :314-317.

61. **Gaur, A.C. (1981).** Phospho microorganism and various transformation in compost technology, Project Field Document No. 13. Rome: FAO.

62. **Gewaily . E . M , Fatma . I . El-zamik, Tomader . T . El-Hadidy , H . I . Abd El-Fattahl and Seham . H . Salem .(2006).** Efficiency of biofertilizers , organic and inorganic amendments application on growth and essential oil of Marjoram (*Majorana hortensis* L.) . Plants grown in sandy and calcareous soil .

63. **Gliessman, S.R., M.R. Werner, S.L. Swezey, E. Caswell, J. Cochran, and F. Rosado-May.(1996).** Conversion to organic strawberry management changes ecological processes. Calif. Agric. 50:24–31.

64. **Gilani. S.S. and Bahmanyar. M.A.(2008).** Impact of Organic Amendments with and Without Mineral Fertilizers on Soil Microbial Respiration Journal of Applied Sciences.8.: 642-647.

65. **Graham, M. H.; and Haynes, R. J.(2005) -** Organic matter accumulation and fertilizer-induced acidification interact to affect soil microbial and enzyme activity on a long-term sugarcane management experiment, Biology and Fertility of Soils, Volume 41(4), 249-256.

66. **Greenwood D.J., Verstraeten L.M.J., Draycott A. and Sutherland R.A. (1987).** Response of winter wheat to N-fertiliser: Dynamic model. *Fertilizer Research* 12:139-156.
67. **Greenwood D.J., Lemaire G., Gosse G., Cruz P., Draycott A. and Neeteson J.J. (1990).** Decline in percentage N of C3 and C4 crops with increasing plant mass. *Annals of Botany*, 66: 425-436.
68. **Greenwood, D.J., Rahn, C.R., Draycott, A.D., Paterson, C.D. and Vaidyanathan, L.v.v. (1996).** Modelling and measurement of the effects of fertilizer-N and residue incorporation on N dynamics in vegetable cropping. *Soil Use and Management*, 12: 13-24.
69. **Gupta, A.P., S.R. Antil and P.R. Narwal, (1988).** Effect of farmyard manure on organic carbon, available N and P contents of soil during different periods of wheat growth. *J. Indian Soil Sci.*, 36: 269–73.
70. **Haiyan .Chu, Takeshi, Fujii, Morimoto. Sho, .Xiangui. Lin, Kazuyuki. Yagi .Junli. Hu and .Jiabao. Zhang .(2006).** Community Structure of Ammonia-Oxidizing Bacteria under Long-Term Application of Mineral Fertilizer and Organic Manure in a Sandy Loam Soil, 73(2): 485–491.
71. **Hamail, A. FL; Helalia, A. M.; Ghaly, F. M. (1996).** Effect of nitrophoska and sagral fertilizers on growth and yield common bean (*Phaseolus vulgaris* L.). *J. Agric. Sci., Mansoura Univ.*, v. 21, n. 1, p. 351-360.
72. **Henriksson .Elisabet and Henriksson .Lars Eric.(1976)** .The Bacteria *Azotobacter*, *Beggiatoa*, and *Desulfovibrio* in the Surtsey soil. Institute of Physiological Botany of the University of Uppsala, S- 75 1 2 1 Uppsala, Sweden.
73. **Herencia.J. F, Ruiz-Porras. J. C, Melero. S, Garcia-Galavis. P. A, Morillo. E and Maqueda. C. (2007).** Comparison between Organic and Mineral Fertilization for Soil Fertility Levels, Crop Macronutrient Concentrations, and Yield. *American Society of Agronomy J* 99:973-983.
74. **Hofman G., Demyttenaere P., de Ossemerct C.I.G. and van Ruymbeke M., (1989).** Nitrate nitrogen losses by leaching in a region of intensive cultivation of vegetables. In "Protection of water quality from harmful emissions with special regard to nitrate and heavy metals".

Proceedings of the 5th International Symposium of CIEC [edited by Welte, E.;Szabolcs, I.], Budapest, Hungary; CIEC, 279-283.

75. **Insam, H. (1990).** Are the Soil Microbial Biomass and Basal Respiration Governed by Climate Regime?, *Soil Biol Biochem*, 22, 525-532.

76. **Isermeyer . H, (1952)**– Eine einfache Methode Zur Bestimmung der Bodenatmung und der carbonate im Boden . *Zpflan Zenernaehr Bodenkd* , Vol 56 , 26 – 38 .

77. **Jacksan M . L (1958)** . Soil chemical analysis . Prentic Hall Inc . Englewood Cliffe . N . J . pp 151- 153 and 331-334 .

78. **Jaggi , W, (1976)** – Die Bestimmung der CO₂ – Bildung als Mab de Boden biologischen Aktivitat . *Schw Landw foresch* , vol 15 , 371-380.

79. **Jeziarska-Tys.Stefania., Frac. Magdalena.(2008).**Effect of sewage sludge on selected Microbiological and Biochemical Indices Of Soil Fertility In View Of Domestic And World Wide Studies 12(2), 393-407.

80. **JECFA.(1995).** “Evaluation of certain food additives and contaminants.” 29-35: World Health Organization.

81. **Jürgens-Gschwind, S.,(1989).** Ground water nitrates in other developed countries (Europe) - relationships to land use patterns. In *Nitrogen Management and Ground Water Protection. Development in Agricultural and Managed Forest Ecology.* (ed. R.F. Follet) 21, pp 75 - 138. Amsterdam: Elsevier.

82. **Kennedy, A.C. (1999).** Bacterial diversity in agroecosystems, *Agriculture, Ecosystems and Environment*, 74, 65–76.

83. **Koji. Nagaya, Kazuaki . Ogino , Tomoko. Shimizu ,Ayumi.Tsunekawa, Kazunuori. Yabemasahiro.Imagawa .(2005).** Effects of Foliar Application of Microelements on Nitrate Reductase Activity and Nitrate Concentration in Cabbage *Research Bulletin of the Aichi-ken Agricultural Research Center* Vol.;NO.37;Page.81-85(2005).

84. **Kolota .Eugeniusz, Adamczewska- Sowińska. Katarzyna.(2007).** Efficiency of Entec 26 - A New Concept Nitrogen Fertilizer in White Head Cabbage Production.vol (69).p:73-80.

85. **Laboski, C.A.M., and J.A. Lamb. (2003).** Changes in soil test phosphorus concentration after application of manure or fertilizer. *Soil Sci. Soc. Am. J.* 67:544–554.

86. **Lawes. J.B, Gilbert. J. H. and Warington. R., . Agric. J. R.(1882).** SOC. *England*, 1882, 18, 1.
87. **Lipman, J.G.; Mclean, H. & Lint, H.C.(1916).** Sulphur oxidation in soils and its effect on the availability of mineral phosphatases. *Soil Sci.*, 1:533-539.
88. **Lombi , E ;Zhao , F ; Fuhrmann, M ;Ma, L.Q and Mcgrath , S.P.(2002).**Arsenic distribution and speciation in the fronds of the hyperaccumulator *pteris vittata* . *New phytologist* 150:195-203.
89. **Lorenz, H.P., Schlaghecken J., Engl G., Maync A., & Zegler J. (1989).** Ordnungsgemäße Stickstoff-Versorgung im Freiland - Gemüsebau - KNS system. Rheinland Phalz: Ministerium Fur Landwirtschaft, Weinbau und Forsten ISSN 0931-9026 1089391-3000.
90. **Lynch, J.M. (1983).** Soil Biotechnology, Microbiological Factors in Crop Productivity. Blackwell Scientific Publications, Oxford.
91. **MAFF .(1994).** Fertilizer Recommendations for agricultural and horticultural crops. Reference Book 209 6th ed. London: HMSO.
92. **Mahmoud, A. R.; Hafez, M. M.; ABD EL-AAL, F. S,(2004).** Comparative study for using organic manure as individual and/or mixing it with chemical fertilizer and their effects on the productivity of *Vicia faba* plants. *J. Agric. Sci., Mansoura Univ.*, v. 29, n. 3, p. 1345-1354.
93. **Mandal A, Patra AK, Singh D, Swarup A, Ebhin Masto R .(2007).** Effect of long-term application of manure and fertilizer on biological and biochemical activities in soil during crop development stages. Division of Soil Science and Agricultural Chemistry, Indian Agricultural Research Institute, New Delhi 110, 012, India. Dec;98(18):3585-92. Epub 2007 Jan 5.
94. **Mandic, L. ; Djukic, D. ; Stevovic V. (2005) -** The Effect of Different Kinds of Fertilizers on the Number of Azotobacters in the Smonitza Type of Soil Under Maize and the Yield of Maize, *Acta Agriculturae Serbica*, Vol X (20), 11-22.
95. **Marinari, S ; G, Masciandro; Ceccanti ; and S, Grego(2000) –** Influence of organic and mineral fertilizer on soil biological and physical properties, *Bioresource Technology*, 72(1), 9-17.
96. **Metcalf, Eddy, 1978,** Wastewater Engineering – Treatment and Reuse, 4 th edition, McGraw Hill.

97. **Monib M., I. Hosny, Y. B. Besada, J. Szegi (1984).** Seed inoculation of castor oil seed plant and its effect on nutrient uptake. *Soil Biology and Conservation of the biosphere* 2; 723-732.
98. **Moreno, J.L., Hernández, T., and Garcia, C. (1999).** Effects of a Cadmium-contaminated Sewage Sludge Compost on Dynamics of Organic Matter and Microbial Activity in an Arid Soil, *Biol Fertil Soils*, 28, 230-237.
99. **Musa, M.M. (1967 ;1974);** Annual Report, Gezira Research Station . Medani Sudan.
100. **Naeem, M., J. Iqbal and M.A.A. Bakhsh, (2006).** Comparative Study of Inorganic Fertilizers and Organic Manures on Yield and Yield Components of Mungbean (*Vigna radiat L.*). *J. Agric. Soc. Sci.*, 2:227–9.
101. **Olsen , R . S . ; C . V . Cole ; F . S . Watanable and L . A . Dean . (1954) .** Estimation of available phoshorus in soil by extraction with sodium bicarbonate . U S D A circular No . 939 .
102. **Parham. J.A.Deng.S.P . DA . H.N. Sun. H .Y. Raun . W. R.(2003) .** Long – term cattle manure in soil and effect on soil microbial populations and community structur.
103. **Parkin, T.B, J.W. Doran, and E. Franco-Vizcaino.(1996).** Field and laboratory tests of soil respiration.p.231-246. In: J.W. Doran and A.J. Jones (eds.) *Methods for assessing soil quality*. Soil Sci.Soc. Am. Spec. Publ. 49. SSSA, Madison, WI.
104. **Parton, W.J., Schimel D.S., Cole, C.V. and Ojima, D.S., (1987).** Analysis of factors controlling soil organic matter levels in great plains grasslands. *Soil Science Society of America Journal* 51: 1173-1179.
105. **Phongran, S.; Mosler A. R. (2003).** Impact of organic residue management on nitrogen use efficiency in an annual rice cropping sequence of lowland central Thailand. *Nutr. Cycl. Agroecosyst.*, v. 66, p. 233-240.
106. **Pikovskaya, R.L (1948).** Mobilization of phosphate in soil in connection with vital activity of some microbial species. *Mikrobiologiya*. 17: 362-370.
107. **Postgate, J. R. (1974).** Evaluation within nitrogen fixation systems. In *evolution in the microbial world*. Ed. M. Carlisle & J. Skehel, symp. Soc. gen. Microbiol., pp. 263-292. Cambridge University Press.

108. **Prochette, P., R.L. Desjardins, and E. Pattey.(1991).** Spatial and temporal variability of soil respiration in agricultural fields. *Can. J. Soil Sci.* 71:189-196.
109. **Quelhas dos Santos, J., (2001),** Fertilização & Ambiente: Reciclagem Agro-Florestal de Resíduos e Efluentes, Publicações Europa-América.
110. **Rahn C.R., Vaidyanathan, L.V. and Paterson, C.D. (1992).** Nitrogen residues from Brassica crops. *Aspects of Applied Biology* 30: 263 - 270.
111. **Rahn C.R., Paterson C.D. and Vaidyanathan L.V.V. (1993).** Improving the use of nitrogen in Brassica rotations. *Acta Horticulturae*, 339 p 207 - 218.
112. **Rahn, C.R., Greenwood, D.J and Draycott, A.(1996).** Prediction of nitrogen fertilizer requirement with HRI WELL_N Computer Model. In: *Progress in Nitrogen Cycling. Proceedings of 8th Nitrogen Fixation Workshop*, University of Ghent, 5-8 September 1994. (eds van Cleemput, O., Hofman, G., and Vermoesen, A.) pp255-258. Dordrecht: Kluwer(A).
113. **Rahn C.R., Paterson C.D., Willison, T., and Lillywhite R. (1996).** The Fate of Nitrogen from Brassica crop residues in intensive cropping rotations. In *Proceedings of COST 814 Conference on 'Nitrogen Supply and Nitrogen Fixation of Crops for Cool and Wet Climates.* September 1995 Tromsø Norway. Published by European Commission Brussels. p 219 - 226 (B).
114. **Rahn C.R., Shepherd M.A. and Hiron R.W.P., (1996).** The effect of water supply on the response of onions and calabres to starter solutions. *Acta Hort., Proc. Workshop Ecological Aspects Veg. Fert.* (eds **Hahndel, R. and Wichman, W.**) *Acta Hort.* 428: 141-150
115. **Rahn C.R., Paterson C.D. and Vaidyanathan L.V.V.(1998).** The use of measurements of soil mineral N in understanding the response of crops to fertilizer nitrogen in intensive cropping rotations. *Journal of Agricultural Science.* In Press.
116. **Rao, N.S.S. (1982)** Biofertilizer in Agriculture, and Agriculture & Forestry. 3rd Ed. Oxford and IBH Publishing Company Pvt. Ltd. New Delhi, India. Pp. 1-14, pp69-82.
117. **Richards L .A . (1962) .** Diagnosis and improvement of saline and alkaline soils . *Agricultural hand book no 60 .* United states Department of agriculture .

118. **Ruiz, J.C.(2002).** Influencia del cambio de un sistema de nutrición mineral o orgánica en las propiedades de suelos hortícolas de regadío. Ph.D. Universidad de Córdoba.
119. **Saad, E. M. (1999).** Effect of some organic manures and microelements on greenhouse cucumber. 141 p. MSc. Thesis, Faculty of Agriculture.
120. **Sady .W, Wojciechowska. R, Rozek. S(1997).** The Effect of form and placement Of N on yield and nitrate content Of white Cabbage. International Society for Horticultural Science .563.
121. **Saha , N . Das – Ac , and D. Mukherjee . (1995) .** Effect of decomposition of organic matter on activities of microorganisms and availability of nitrogen . phosphorus , and sulphur in soil . Journal of the Indian society of soil science 43:2,210-215 .
122. **Sanyal, S.K., and S.K. De Datta. (1991).** Chemistry of phosphorus transformations in soil. Adv. Soil Sci. 16:1–120.
123. **Scaife M.A. and Turner M.K., (1985).** Effect of nitrogen top-dressing on Brussels sprout flavour. Tests of Agrochemicals and Cultivars, 106: 172-173.
124. **Scaife M.A. and Wurr D.C.E.,(1990).** Effects of nitrogen and irrigation on hollow stem of cauliflower (Brassica oleracea var Botrytis). J. Hort. Sci., 65: 25-29
125. **Scaife M. A. and Bar-Yosef, B. (1995).** Nutrient and fertilizer management in field grown vegetables . In IPI - Bulletin No 13. International Potash Institute Basel Switzerland.
126. **Shkenan varyou.N.M.(1991).**Reduce the negative impact of nitrogen fertilizers ,agriculture chemistrization ,No6 .in Russian.
127. **Singh S., K. K. Kapoor (1999).** Inoculation with phosphate – solubilizing microorganisms and a vesicular –arbuscular mycorrhizal fungus improves dry matter yield and nutrient uptake by wheat grown in a sandy soil. Biol. Fertil. Soils 28,139- 144.
128. **Sinaj, S.; Traore, O.; Frossard, E.(2002).** Effect of compost and soil properties on the availability of compost phosphate for white clover (*Trifolium repens* L.). Nutr. Cycl. Agroecosyst., v. 62, p. 89-102.
129. **Sims, J.T. (1995).** Organic wastes as alternative nitrogen sources. In P.E. Bacon (ed.) Nitrogen fertilization in the environment. Marcel Dekker, New York.

130. **Sørensen J.N.,(1996).** Improved N efficiency in vegetable production by fertilizer placement and irrigation. *Acta Hort., Proc. Workshop Ecological Aspects Veg. Fert.* (eds Hahndel, R. and Wichman, W.)*Acta Hort.* 428: 131-140.
131. **Srivastava , M; MA,L.Q;Sing , N.and Sing , S.(2005).**Antioxidant responses of hyper-accumulator and sensitive fern species to arsenic . *J .EXP.Bot .* 56.1335-1342.
132. **Stewart, M.W., W.D. Dobb, E.A. Johnston and J.T. Smyth, (2005).** The Contribution of Commercial Fertilizer Nutrients to Food Production.*Agron. J.*, 97: 1–6.
133. **Stone, D.A. and Rowse, H.R., (1992).** Reducing the nitrogen requirement of vegetable crops by precision fertilizer injection. *Aspects of Applied Biology* 30:399-402.
134. **Subba Rao, N.S. (1994).** Mikroorganisme Tanah dan Pertumbuhan Tanaman.Jakarta: Penerbit Universitas Indonesia.
135. **Sundara Rao, W.V.B. and M.K. Sinha.(1963).** Phosphate dissolving microorganism in the soil and rhizosphere. *Indian Journal of Agricultural Science* 33: 272-278.
136. **Suresh, K.D., G. Sneh, K.K. Krishn and C.M. Mool,(2004).** Microbial biomass carbon and microbial activities of soils receiving chemical fertilizers and organic amendments. *Archives Agron. Soil Sci.*, 50:641–7.
137. **Taiz,L. and Zeiger , E . (2002):** Plant physiology . Sunderland , MA: Sinauer . 690 pp.
138. **Theodora, M., S.L. Anastasios and A.G. Athanasios, (2003).** Effects of injected liquid cattle manure on growth and yield of winter wheat and soil characteristics. *Agron. J.*, 95: 592–60.
139. **Tisdale, S.L., W.L. Nelson, and J.D. Beaton. (1985).** Soil fertility and fertilizers. 4th ed. Macmillan Publ. Co., New York.
140. **Van Cleue, Keith ; A.Moore, Terry (1978) -** Cumulative effects of nitrogen, phosphorus, and potassium fertilizer additions on soil respiration, pH, and organic matter content, *Soil Science Society of American Journal*, vol 42,121-124.
141. **Valarini, P.J.; Frighetto, R.T.A.; Tokeshi, H. &Oliveira, D.A. (1997).** Action of effective microorganisms (EM) in soil of intensive agricultural system in Brazil. In: *International Conference On Kyusei*

Nature Farming, 5., Bangkok, 1997. Proceedings angkok, Asia Pacific Natural Agriculture Network, p.15-16.

142. **Van den Boogaard R. and Thorup-Kristensen, K., (1997).** Effects of nitrogen fertilization on growth parameters and soil nitrogen depletion in cauliflower. *Acta Agric Scand., Sect. B, Soil and Plant Sci.* In press

143. **Van Veen, J. A. and Frissel, M.J., (1981).** Simulation model of the behaviour of N in soil. In: *Simulation of nitrogen behaviour of soil-plant systems.* (eds Frissel M.J. and van Veen, J.A.) pp126-144. Wageningen: Centre for Agricultural Publishing and Documentation.

144. **Vikram , A ; Alagawadi , R .A ; Hamzehzarghani,H and U.P.Krishnaraj.(2007).** Factors Related to the Occurrence of Phosphate Solubilizing Bacteria and their Isolation in Vertisols . *International Journal of Agriculture Research* 2(7) ; 571-580 .

145. **Vineela, C ; Wani, S.P ; Srinivasarao, Ch ; Padmaja, B ; and Vittal, K.P.R (2008)** – Microbial properties of soils as affected by cropping and nutrient management practices in several long-term manural experiments in the semi- arid tropics of India, *Applied Soil Biology*,vol 40(1), 165-173.

Volk, T.(1994). The soil's breath. *Natural History* November/94. Background & Interpretive Guide for Individual Tests.

146. **Walinga . I ., J . J Van Der Lee ., V . J . G Houba . W.Van Vark . I . Novozamsky , (1995) .** *Plant Analysis Manual . Kluwer Academic Publishers . London .*

147. **Weier, U.,(1992).** Effect of splitting N fertilizer on yield of Broccoli. In: *Versuche in Deutschen Gartenbau* 1992, No. 26. Bonn.

148. **Whiting , S.N.(2001).** Rhizosphere bacteria mobilize zinc for hyperaccumulation by *thlaspi caerulenscens* . *Environ . SCI . Technol .* 35 : 3144-3150.

149. **Wilkinson, S.R., (1979).** Plant nutrient and economic value of animal manures. *J. Animal Sci.*, 48: 121–33.

150. **Wintz, H . Fox, T and Vulpe , C. (2002).**Functional Genomic and Gene Regulation in Biometals Research : Responses of plants to iron , Zinc and Copper deficiencies . *Biochemical society Transaction .* 30(4): 766-768.

151. **Wong, J.W.C., K.K. Ma, K.M. Fang and C. Cheung, (1999).** Utilization of manure compost for organic farming in Hong Kong. *Bio-resource Technol.*, 67: 43–6.
152. **Yang, Lan-fang ; Cai, Zu-cong (2006)** – Soil Respiration During a Soybean Growing Season, *Pedo sphere*, vol 16(2), 192-200.
153. **Zane, F.L. and D.D. Basil, (1980).** Residual effects of dairy cattle manure on plant growth and soil properties. *Agron. J.*, 72: 123–30.
154. **Zeidan, M.; Kabesh, M. O.; Saber, M. S. M.(2001).** Utilisation of biofertilizers in field crop production. 14- Effect of organic manuring and biofertilisation on yield composition of two faba bean varieties in a newly reclaimed soil. *Egypt. J. Agron.*, v. 23, n. 1, p. 47-54.
155. **Zhang Qi- chan, Wang Guang- huo, (2006)** -__The effect of different fertilization treatments on ecological characteristics of microorganisms in paddy soil, Department of natural resources , Zhejiang university science pp376-380.
156. **Zhong .Wenhui , Gu. Ting , Wang. Wei , Zhang. Bin , Lin. Xiangui , Huang. Qianru and Shen. Weishou .(2008).**The effects of mineral fertilizer and organic manure on soil microbial community and diversity. *Plant and Soil*.

The Effect of Organic Fertilization in Soil Microorganisms and the Productivity of Two Crops of Brassica Family (Cabbage - Cauliflower)

Abstract :

An experiment was carried out to study the effect of organic fertilization and nitrogen fertilizer on cabbage and cauliflower crops for two seasons, the included 6 treatments (control , cow manure , sheep manure , chicken manure , compost , and nitrogen fertilizers). Organic fertilizers were added as similar rate of N in urea (50 kg/N/ha) according to the content of soil nitrogen and Ministry of Agric. and Agra. Reform.S.(Recommendation) .

This study has been done on some biological properties of soil (the number of some physiological groups of soil microorganisms , soil respiration ,in addition to nitrate concentration in plant and (Cabbage – Cauliflower) productivity were determined.

This investigation shows for Cabbage crop that the number of ,Bacterial assimilation mineral form of nitrogen, Actinomycetes , *Azotobacter*, Phosphate solubilizing microorganisms, Mesophilic cellulose decomposing microorganisms and Spor-forming bacteria (*Bacillus*) increased in most treatments amended with organic fertilizers comprising to control , whereas the number of, Actinomycetes , *Azotobacter* , Phosphate solubilizing microorganisms Fungi , Mesophilic cellulose decomposing microorganisms and *Clostridium pasteurianum* decreased in treatment amended with nitrogen fertilizers as compared to control.

No significant differences noticed in the intensity of soil respiration test between treatments at two seasons for cabbage crop.

The concentration of nitrate increased on the heads of Cabbage in treatments amended with nitrogen fertilizers and treatment amended with cow manure at the first season ,whereas the increasing of nitrate on the head of Cabbage at second season was in treatment amended with chicken manure.

The yield of cabbage increased in treatments amended with chicken manure at the first season, whereas the productivity of Cabbage increased in treatments amended with cow manure at the second season.

The results show on Cauliflower crop that the number of Bacterial assimilation mineral form of nitrogen , Actinomycetes , *Azotobacter*, Phosphate solubilizing microorganisms , Mesophilic cellulose decomposing microorganisms and Spor-forming bacteria (*Bacillus*) in most treatments that amended with organic fertilizers was increasing in compared to control , and the number of Ammonification bacteria , Actinomycetes , *Azotobacter* , Phosphate solubilizing microorganisms, Fungi , Mesophilic cellulose decomposing microorganisms and *Clostridium pasteurianum* in the treatment that amended with nitrogen fertilizers was decreasing ,this treatment was the lowest in compared to all treatments even the control.

No significant differences noticed in the intensity of soil respiration test between treatments at first seasons for Cauliflower crop. , whereas the intensity of soil respiration increased in treatments amended with nitrogen fertilizer at the second season .

The concentration of nitrate increased on the floral curd of Cauliflower in treatments amended with nitrogen fertilizers and treatment amended with cow manure at the first season ,whereas the increasing of nitrate on the floral curd of Cauliflower at second season was in treatment amended with cow manure.

The yield of Cauliflower increased in treatments amended with chicken manure at the first season, whereas the productivity of Cauliflower increased in treatments amended with cow manure at second season.

Key words: soil , Organic Fertilization ,Nitrogen fertilizer, Microorganisms , Soil respiration ,nitrate , Productivity .
